

Thursday, July 8, 2010

Problem 4. Let P be a point inside the triangle ABC . The lines AP , BP and CP intersect the circumcircle Γ of triangle ABC again at the points K , L and M respectively. The tangent to Γ at C intersects the line AB at S . Suppose that $SC = SP$. Prove that $MK = ML$.

Problem 5. In each of six boxes $B_1, B_2, B_3, B_4, B_5, B_6$ there is initially one coin. There are two types of operation allowed:

Type 1: Choose a nonempty box B_j with $1 \leq j \leq 5$. Remove one coin from B_j and add two coins to B_{j+1} .

Type 2: Choose a nonempty box B_k with $1 \leq k \leq 4$. Remove one coin from B_k and exchange the contents of (possibly empty) boxes B_{k+1} and B_{k+2} .

Determine whether there is a finite sequence of such operations that results in boxes B_1, B_2, B_3, B_4, B_5 being empty and box B_6 containing exactly $2010^{2010^{2010}}$ coins. (Note that $a^{b^c} = a^{(b^c)}$.)

Problem 6. Let a_1, a_2, a_3, \dots be a sequence of positive real numbers. Suppose that for some positive integer s , we have

$$a_n = \max\{a_k + a_{n-k} \mid 1 \leq k \leq n-1\}$$

for all $n > s$. Prove that there exist positive integers ℓ and N , with $\ell \leq s$ and such that $a_n = a_\ell + a_{n-\ell}$ for all $n \geq N$.