

## 20th USA Mathematical Olympiad

April 23, 1991

Time Limit:  $3\frac{1}{2}$  hours

1. In triangle  $ABC$ , angle  $A$  is twice angle  $B$ , angle  $C$  is obtuse, and the three side lengths  $a, b, c$  are integers. Determine, with proof, the minimum possible perimeter.
2. For any nonempty set  $S$  of numbers, let  $\sigma(S)$  and  $\pi(S)$  denote the sum and product, respectively, of the elements of  $S$ . Prove that

$$\sum \frac{\sigma(S)}{\pi(S)} = (n^2 + 2n) - \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}\right)(n+1),$$

where “ $\Sigma$ ” denotes a sum involving all nonempty subsets  $S$  of  $\{1, 2, 3, \dots, n\}$ .

3. Show that, for any fixed integer  $n \geq 1$ , the sequence

$$2, 2^2, 2^{2^2}, 2^{2^{2^2}}, \dots \pmod{n}$$

is eventually constant.

[The tower of exponents is defined by  $a_1 = 2$ ,  $a_{i+1} = 2^{a_i}$ . Also  $a_i \pmod{n}$  means the remainder which results from dividing  $a_i$  by  $n$ .]

4. Let  $a = (m^{m+1} + n^{n+1})/(m^m + n^n)$ , where  $m$  and  $n$  are positive integers. Prove that  $a^m + a^n \geq m^m + n^n$ .

[You may wish to analyze the ratio  $(a^N - N^N)/(a - N)$ , for real  $a \geq 0$  and integer  $N \geq 1$ .]

5. Let  $D$  be an arbitrary point on side  $AB$  of a given triangle  $ABC$ , and let  $E$  be the interior point where  $CD$  intersects the external common tangent to the incircles of triangles  $ACD$  and  $BCD$ . As  $D$  assumes all positions between  $A$  and  $B$ , prove that the point  $E$  traces the arc of a circle.

