

**23<sup>rd</sup> United States of America Mathematical Olympiad**

**April 28, 1994**

**Time Limit:  $3\frac{1}{2}$  hours**

1. Let  $k_1 < k_2 < k_3 < \cdots$  be positive integers, no two consecutive, and let  $s_m = k_1 + k_2 + \cdots + k_m$  for  $m = 1, 2, 3, \dots$ . Prove that, for each positive integer  $n$ , the interval  $[s_n, s_{n+1})$  contains at least one perfect square.
2. The sides of a 99-gon are initially colored so that consecutive sides are red, blue, red, blue,  $\dots$ , red, blue, yellow. We make a sequence of modifications in the coloring, changing the color of one side at a time to one of the three given colors (red, blue, yellow), under the constraint that no two adjacent sides may be the same color. By making a sequence of such modifications, is it possible to arrive at the coloring in which consecutive sides are red, blue, red, blue, red, blue,  $\dots$ , red, yellow, blue?
3. A convex hexagon  $ABCDEF$  is inscribed in a circle such that  $AB = CD = EF$  and diagonals  $AD$ ,  $BE$ , and  $CF$  are concurrent. Let  $P$  be the intersection of  $AD$  and  $CE$ . Prove that  $CP/PE = (AC/CE)^2$ .
4. Let  $a_1, a_2, a_3, \dots$  be a sequence of positive real numbers satisfying  $\sum_{j=1}^n a_j \geq \sqrt{n}$  for all  $n \geq 1$ . Prove that, for all  $n \geq 1$ ,

$$\sum_{j=1}^n a_j^2 > \frac{1}{4} \left( 1 + \frac{1}{2} + \cdots + \frac{1}{n} \right).$$

5. Let  $|U|$ ,  $\sigma(U)$  and  $\pi(U)$  denote the number of elements, the sum, and the product, respectively, of a finite set  $U$  of positive integers. (If  $U$  is the empty set,  $|U| = 0$ ,  $\sigma(U) = 0$ ,  $\pi(U) = 1$ .) Let  $S$  be a finite set of positive integers. As usual, let  $\binom{n}{k}$  denote  $\frac{n!}{k!(n-k)!}$ . Prove that

$$\sum_{U \subseteq S} (-1)^{|U|} \binom{m - \sigma(U)}{|S|} = \pi(S)$$

for all integers  $m \geq \sigma(S)$ .